

Twist-3 pion and kaon distribution amplitudes from the instanton vacuum with flavor SU(3) symmetry breaking

Seung-il Nam^{*} and Hyun-Chul Kim[†]

*Department of Physics and Nuclear Physics & Radiation Technology Institute (NuRI),
Pusan National University, Busan 609-735, Republic of Korea*

Abstract

We investigate the twist-3 pion and kaon distribution amplitudes of the pseudoscalar ($\phi_{\pi,K}^p$) and pseudotensor ($\phi_{\pi,K}^\sigma$) types, based on the effective chiral action from the instanton vacuum. Flavor SU(3) symmetry breaking effects are explicitly taken into account. The Gegenbauer moments and the moments of the distribution amplitudes ($\langle \xi^m \rangle$) are also computed. Our results are summarized as follows: $a_{2,\pi}^p \sim 0.4$, $a_{1,K}^p \sim 0.02$ and $a_{2,K}^p \sim 0.14$, and $a_{2,\pi}^\sigma \sim 0.02$ and $a_{1,K}^\sigma \sim a_{2,K}^\sigma \sim 0$; $\langle \xi^2 \rangle_\pi^p \sim \langle \xi^2 \rangle_K^p \sim 0.37$ and $\langle \xi \rangle_K^p \sim 0$, and $\langle \xi^2 \rangle_\pi^\sigma \sim \langle \xi^2 \rangle_K^\sigma \sim 0.20$ and $\langle \xi \rangle_K^\sigma \sim 0$. We compare our results with those from the QCD sum rules. We also discuss the relevant Wilson coefficients which were analyzed recently in chiral perturbation theory.

PACS numbers: 11.15.Tk, 14.40.Aq

Keywords: Twist-3 pion and kaon distribution amplitudes, Nonlocal chiral quark model, Instanton vacuum

^{*}Electronic address: sinam@pusan.ac.kr

[†]Electronic address: hchkim@pusan.ac.kr

I. INTRODUCTION

The meson light-cone distribution amplitude (DA) plays an essential role in describing exclusive hadronic reactions [1, 2, 3, 4]. While the leading-twist DAs provide major contribution to those processes, higher-twist DAs are suppressed by the inverse of the momentum transfer Q^2 , so that higher-twist DAs are less significant in studying high-energy exclusive processes. Because of this fact, not many investigations on higher-twist DAs were performed [5, 6, 7, 8, 9, 10, 11], compared to the leading-twist ones. However, higher-twist DAs should be still important, since they give crucial information on the soft part of exclusive hadron reactions, in particular, in the smaller Q^2 region. Moreover, it describes the transverse motion of quarks inside the meson [6, 11].

In the present work, we aim at investigating the two-particle twist-3 pion and kaon DAs within the framework of the nonlocal chiral quark model (χ QM) from the instanton vacuum, taking into account SU(3) symmetry breaking effects. The instanton model of the QCD vacuum provides a good framework to study the structure of light mesons, since spontaneous chiral symmetry breaking is well realized in the instanton vacuum via the quark zero mode [12, 13, 14]. This instanton vacuum model was later extended by introducing the current-quark masses [15, 16, 17]. It was assumed in the model that the large N_c expansion is the reasonable one and the results were obtained in the leading order in this expansion. In the present approach, we employ the modified effective chiral action with flavor SU(3) symmetry breaking effects derived from the instanton vacuum [15, 16, 17].

The model has been applied to describe the leading-twist meson DAs [18, 19, 20, 21, 22]. One of the two-particle twist-3 pion DAs was already studied within the nonlocal χ QM in the chiral limit [21]. There are two independent particle-two twist-3 distribution amplitudes defined as:

$$\phi_{\mathcal{M}}^p(u) = \frac{\sqrt{2}(P \cdot \hat{n})(m_f + m_g)}{m_{\mathcal{M}}^2 F_{\mathcal{M}}} \int_{-\infty}^{\infty} \frac{d\tau}{\pi} e^{-i(2u-1)\tau P \cdot \hat{n}} \langle 0 | \bar{\psi}_f(\tau \hat{n}) i \gamma_5 \psi_g(-\tau \hat{n}) | \mathcal{M}(P) \rangle, \quad (1)$$

$$\begin{aligned} \phi_{\mathcal{M}}^\sigma(u) = & -\frac{6\sqrt{2}(m_f + m_g)}{m_{\mathcal{M}}^2 F_{\mathcal{M}}} \int_{-\infty}^{\infty} \frac{d\tau}{\pi} \int_0^u dv e^{-i(2v-1)\tau P \cdot \hat{n}} \\ & \times \langle 0 | \bar{\psi}_f(\tau \hat{n}) i (P \not{\hat{n}} - P \cdot \hat{n}) \gamma_5 \psi_g(-\tau \hat{n}) | \mathcal{M}(P) \rangle, \end{aligned} \quad (2)$$

where the subscript $\mathcal{M}$ denotes either the pion ($\mathcal{M} = \pi$) or the kaon ($\mathcal{M} = K$) with on-mass-shell momentum P in the light-cone frame. u and n^μ stand for the longitudinal momentum fraction and the light-like vector satisfying $n^2 = 0$, respectively. The spatial separation between the quarks inside the meson is represented by $\tau \cdot n$. In the present work, we drop the Wilson line by considering the light-cone gauge, $A \cdot n = 0$. m_f and m_g are the corresponding current quark masses with given flavors f and g . As for the present case, we assign them as $(f, g, \mathcal{M}) = (s, u, K^+)$ and (d, u, π^+) . $m_{\mathcal{M}}$ denotes the corresponding meson mass. $F_{\mathcal{M}}$ is the corresponding meson decay constant which is the normalization constant for the DAs. The DAs of Eqs. (1) and (2) satisfy the following normalization conditions:

$$\int_0^1 du \phi_{\mathcal{M}}^p(u) = \int_0^1 du \phi_{\mathcal{M}}^\sigma(u) = 1. \quad (3)$$

The present work is organized as follows: Section II is devoted to the general formalism. The numerical results and discussions are given in Section III. We summarize and draw conclusions in Section IV.

II. GENERAL FORMALISM

In order to evaluate the nonlocal hadronic matrix elements of Eqs. (1) and (2), we start from the low-energy effective QCD partition function derived from the instanton vacuum in Euclidean space as follows [14, 15, 16, 17]:

$$\mathcal{Z} = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} \mathcal{D}\mathcal{M}^a \exp \int d^4x \left[i\bar{\psi}_f^\dagger(x)(i\not{\partial} + im_f)\psi_f(x) - \int \frac{d^4k d^4p}{(2\pi)^8} e^{i(k-p)\cdot x} \sqrt{M_f(k)M_g(p)} \bar{\psi}_f(k) U_{fg}^{\gamma_5} \psi_g(p) \right], \quad (4)$$

where ψ and $\mathcal{M}$ are the fields for the quark and pseudoscalar meson. $M_f(k)$ is the dynamically generated quark mass discussed later. The background pseudoscalar meson field U^{γ_5} is given by

$$U^{\gamma_5} = U(x) \frac{1 + \gamma_5}{2} + U^\dagger(x) \frac{1 - \gamma_5}{2} = 1 + \frac{i}{F_{\mathcal{M}}} \gamma_5 \mathcal{M}^a \lambda^a - \frac{1}{2F_{\mathcal{M}}^2} (\mathcal{M}^a)^2 \dots \quad (5)$$

λ^a is the well-known SU(3) Gell-Mann matrices. Now we are in a position to construct the quark propagator in the instanton ensemble. There have been several approaches for this purpose [15, 16, 17]. In the present work, we make use of the prescription suggested by Pobylitsa [23] and Musakhanov [16]. We attempt to discuss it briefly in order. First, Diakonov *et al.* introduced the quark propagator under the instanton effects by taking into account the following assumption [24]:

$$\langle x | \frac{1}{i\not{\partial} + \not{A}_I + im_F} | x \rangle \simeq \langle x | \frac{1}{i\not{\partial} + im_F} | x \rangle + \frac{\psi_I(x) \psi_I^\dagger(x)}{im_f}, \quad (6)$$

where A_I and $\psi(x)$ stand for the single instanton field and the quark zero mode, respectively. However, this assumption loses information on higher quark loops although it turns out that it works very well from phenomenological point of view. On the contrary, Pobylitsa expand the quark propagator $\langle x | (i\not{\partial} + \not{A}_I + im)^{-1} | x \rangle$ straightforwardly by virtue of the large N_c limit [23]. Thus, all non-planar diagrams are removed from the expansion. One then obtains an integral equation as follows:

$$\begin{aligned} & \langle x | \frac{1}{i\not{\partial} + \not{A}_I + im} | x \rangle \\ &= \langle x | \frac{1}{i\not{\partial} + im_f} | x \rangle + \frac{N}{2VN_c} \text{tr}_c \left[\int d^4Z_I \left[\langle x | (i\not{\partial} + \not{A}_I + im_f) | x \rangle - \frac{1}{\not{A}} \right]^{-1} + (I \rightarrow \bar{I}) \right], \end{aligned} \quad (7)$$

where $Z_{I,\bar{I}}$ indicates the instanton coordinate for the instanton and antiinstanton. By solving Eq. (7) using an ansatz $\sigma(k)$ for the full propagator, $[i\not{\partial} + im_f + \sigma(i\not{\partial})]^{-1}$, one can obtain the current-quark mass dependent dynamical quark mass [16, 23]. Final expression for it can be written as follows:

$$M_f(k) = M_0 F^2(k\bar{\rho}) \left[\sqrt{1 + \frac{m_f^2}{d^2}} - \frac{m_f}{d} \right], \quad \text{where } d = \sqrt{\frac{0.08385}{2N_c}} \frac{8\pi\bar{\rho}}{R^2} \simeq 0.198 \text{ GeV}. \quad (8)$$

We notice that, as a consequence, all the effects from the current-quark mass are included in the bracket of Eq. (8). Here, we employ the standard values for the instanton ensemble, $N/V = 200^4 \text{ MeV}^4$ and $\bar{\rho} \sim 1/3 \text{ fm} \simeq 1/600 \text{ MeV}^{-1}$ for the numerical calculations. From these values, we obtain $M_0 = 0.350 \text{ GeV}$. Although the form factor $F(k)$ can be derived from the Fourier transform of the quark zero mode, we will use the following simple parameterization for it:

$$M_f(k) = M_0 \left[\frac{n\Lambda^2}{(n\Lambda^2 - k^2 + i\epsilon)} \right]^{2n} \left[\sqrt{1 + \frac{m_f^2}{d^2}} - \frac{m_f}{d} \right]. \quad (9)$$

Note that we are now working in Minkowski space, in which the DA are well defined, rather than in Euclidean one. In the present work as done in our previous work [22], we assume that the partition function of Eq. (4) can be analytically continued to Minkowski space. Although we do not provide a firm theoretical proof for this assumption, it turns out that it works qualitatively well from the phenomenological point of view [19, 20, 21]. Λ is the cutoff mass which can be taken as the scale parameter of the present work. We use $\Lambda \simeq 1.0 \text{ GeV}$ for the numerical calculation as in [22]. In order to check the effect of the parameterization of the form factor, we take three different values of the power in Eq. (9), namely, $n = 1, 2$ and 3 . With the scheme discussed above, we can write the expressions for the two-particle twist-3 pion and kaon DAs, $\phi_{\mathcal{M}}^p$ and $\phi_{\mathcal{M}}^\sigma$:

$$\begin{aligned} \phi_{\mathcal{M}}^p(u) = & -\frac{iN_c P_+(m_f + m_g)}{m_{\mathcal{M}}^2 F_{\mathcal{M}}^2} \int \frac{dk_+ dk_- dk_T^2}{(2\pi)^4} \delta[uP_+ - k_+] \\ & \times \text{tr}_d \left[\frac{\sqrt{M_f(k)}}{D_f} \gamma_5 \frac{\sqrt{M_g(k-P)}}{D_g} \gamma_5 \right], \end{aligned} \quad (10)$$

$$\begin{aligned} \phi_{\mathcal{M}}^\sigma(u) = & \frac{6iN_c(m_f + m_g)}{m_{\mathcal{M}}^2 F_{\mathcal{M}}^2} \int_0^u dv \int \frac{dk_- dk_+ d^2 k_T}{(2\pi)^4} \delta[vP_+ - k_+] \\ & \times \text{tr}_d \left[\frac{\sqrt{M_f(k)}}{D_f} (P\not{k} - P_+) \gamma_5 \frac{\sqrt{M_g(k-P)}}{D_g} \gamma_5 \right], \end{aligned} \quad (11)$$

where $P_+ = P \cdot \hat{n}$. D_f denotes the inverse quark propagator, $\not{k} + m_f + M_f(k)$. The trace tr_d denotes the trace over Dirac space. The explicit evaluation of the DAs in Eqs. (10) and (11) is given in Appendix. We set $m_\pi = 140 \text{ MeV}$ and $m_K = 495 \text{ MeV}$ for numerical input. We assume isospin symmetry with $m_u = m_d = 5 \text{ MeV}$. We choose $m_s = 150 \text{ MeV}$ for the strange current quark mass.

III. NUMERICAL RESULTS

In the present Section, we provide the numerical results of the twist-3 pion and kaon DAs. We fix the parameters, using the normalization conditions in Eq. (3). The method is given in Refs. [22] in detail.

In Fig. 1, we depict the pseudoscalar-type DAs ($\phi_{\mathcal{M}}^p$) for the pion (left) and kaon (right). For comparison, we also show the asymptotic DA ($\phi_{\text{Asym.}}^p(u) = 1$) and those of Ref. [11] in each panel. Note that the results of Ref. [11] using the QCD sum rules (QCDSR) were derived at the renormalization scale $\mu = 1 \text{ GeV}$, which is rather compatible to our case. As




FIG. 1: The results of the pseudoscalar twist-3 pion distribution amplitudes ϕ_π^p (left) and kaon ones ϕ_K^p (right). The dot-dashed curves indicate the results of the QCDSR in Ref. [11]. The asymptotic one $\phi_{\text{Asym}}^p = 1$ is drawn in the long-dashed line.

expected from isospin symmetry, the pion DAs are all symmetric as shown in the left panel of Fig. 1. Examining the dependence of the DAs on the power n in Eq. (9), we find a very interesting behavior of the DAs with power n . With $n = 1$ used, the pion DA does not vanish at the end points of u and turns out to be very similar to that of Ref. [11], while the pion DAs are suppressed in the vicinity of the end points and eventually vanish at the end points in the case of $n = 2$ and 3. In this case, the present results look similar to those of Ref. [7]. The reason lies in the fact that for the parameterization of the momentum-dependent quark mass with $n = 1$ there exist nonvanishing terms at the end points after integration over k_- variables.

In the right panel, we depict the kaon DAs, $\phi_K^p(u)$. We see that the kaon DAs are almost symmetric in spite of the mass difference between the light and strangeness quarks. Note that this behavior was already seen even for the leading twist kaon DA in our previous work [22]. It turns out that the present results of the ϕ_K^p are very different in shape, compared to that of Ref. [11]. We conclude that, in our model framework, the pion and kaon DAs show negligible difference.

In Fig. 2, we draw the present results of the pseudotensor twist-3 pion and kaon DAs ($\phi_{\mathcal{M}}^\sigma$) whose asymptotic form is $\phi_{\text{Asym}}^\sigma = 6u(1-u)$. When $n = 1$ is used, the ϕ_π^σ turns out to be almost the same as the asymptotic one. However, as n increases, the ϕ_π^σ are suppressed rather strongly near the end points, whereas they get narrower and larger in the neighborhood of the center ($u = 0.5$). It can be easily understood from the normalization condition for ϕ_π^σ in Eq. (3). Compared to the result of Ref. [11] again, the behavior of the present ϕ_π^σ is of great difference from it. While the present ϕ_π^σ are humped near the center, that of Ref. [11] is rather flat in that region. The pseudotensor twist-3 kaon DAs ϕ_K^σ look similar to the pion ones, as indicated in the right panel of Fig. 2, though they are negligibly asymmetric to the right. However, result of Ref. [11] is shifted to the right and slightly asymmetric.

In order to analyze the $\phi_{\mathcal{M}}^p$ and $\phi_{\mathcal{M}}^\sigma$ in detail, we compute their Gegenbauer moments. By doing that, we can immediately check how much the DAs are deviated from the asymptotic ones. In addition, they give information on the strength of flavor SU(3) symmetry breaking effects for the kaon DAs. The twist-3 DAs can be expanded in terms of the Gegenbauer




FIG. 2: The results of the pseudotensor twist-3 pion distribution amplitudes ϕ_π^σ (left) and kaon ones ϕ_K^σ (right). The dot-dashed curves indicate the results of the QCDSR in Ref. [11]. The asymptotic one $\phi_{\text{Asym}}^\sigma = 6u(1-u)$ is drawn in the dashed line.

polynomials:

$$\phi_{\mathcal{M}}^p(u) = \sum_{m=0}^{\infty} a_{m,\mathcal{M}}^p C_m^{1/2}(\xi), \quad (12)$$

$$\phi_{\mathcal{M}}^\sigma(u) = 6u(1-u) \sum_{m=0}^{\infty} a_{m,\mathcal{M}}^\sigma C_m^{3/2}(\xi), \quad (13)$$

where $\xi = 2u-1$. The orthogonal condition for the Gegenbauer polynomials, the Gegenbauer moments for the $\phi_{\mathcal{M}}^p$ and $\phi_{\mathcal{M}}^\sigma$ can be derived as follows:

$$a_{m,\mathcal{M}}^p = (2m+1) \int_0^1 du C_m^{1/2}(\xi) \phi_{\mathcal{M}}^p(u), \quad (14)$$

$$a_{m,\mathcal{M}}^\sigma = \frac{4m+6}{3m^2+9m+6} \int_0^1 du C_m^{3/2}(\xi) \phi_{\mathcal{M}}^\sigma(u), \quad (15)$$

where m denotes an order of the Gegenbauer moment. The results of the Gegenbauer moments are listed in Table I. First, we consider the pseudoscalar ones, $a_{\mathcal{M}}^p$. As expected from isospin symmetry, it turns out that all odd Gegenbauer moments vanish in the case of the pion DAs. Moreover, we see that the Gegenbauer moments are all negative except for the case of $n=1$. The result of $a_{2,\pi}^p$ is very similar in general to that of Ref. [11]. However, the values of $a_{4,\pi}^p$ and $a_{6,\pi}^p$ are positive in the present work but they are negative in Ref. [11], though in Fig. 1 they are seemingly almost the same each other.

As for the kaon DA, we find that the situation becomes much more interesting. The first Gegenbauer moment $a_{1,K}^p$, which encodes the strength of flavor SU(3) symmetry breaking effects, turns out to be all positive in our calculation. Its value is about 0.02. The smallness of the Gegenbauer moments explains why the kaon DA becomes almost symmetric and rather flat. Note that the even Gegenbauer moments are much smaller than those of the pion DAs. The results of the present work for the kaon DA are much smaller than those of the QCDSR [11].

We now consider the Gegenbauer moments for the $\phi_{\mathcal{M}}^\sigma$. The overall tendency for the sign of the Gegenbauer moments is very similar to that for the $\phi_{\mathcal{M}}^p$. We find that the present

results of the $a_{2\pi}^\sigma$ are much smaller than those of Ref. [6, 11]. Larger values of the $a_{2,\pi}^\sigma$ imply that the DA becomes flat, as shown in Fig. 2. Since we obtain all negative $a_{2,K}^\sigma$, the kaon DAs have narrower shapes. Note that all odd kaon Gegenbauer moments are very small ($\sim 10^{-2}$). Once again, we see the reduced SU(3) symmetry breaking effects.

It is also of great interest to study the moments of the DAs defined as follows:

$$\langle \xi^m \rangle_{\mathcal{M}}^p = \int_0^1 du \xi^m \phi_{\mathcal{M}}^p(u), \quad (16)$$

$$\langle \xi^m \rangle_{\mathcal{M}}^\sigma = \int_0^1 du \xi^m \phi_{\mathcal{M}}^\sigma(u). \quad (17)$$

The results are listed in Table II, being compared with the QCDSR results from Refs. [7, 8, 11]. We observe that $\langle \xi^{2,4,6} \rangle_\pi^p$ are comparable with the QCDSR results [7, 11]. We find that the values of $\langle \xi^{1,3,5} \rangle_K^p$ are small in comparison with Refs. [7, 11], although the even moments are still similar. The results of Ref. [8] are slightly larger than the present ones in general.

As for the moments of the $\phi_{\mathcal{M}}^\sigma$, it turns out that the present results are comparable with those of Ref. [11], though the difference between them is getting larger for higher moments. The situation is similar when we compare our results with those of Ref. [8]. We note that the odd moments become all negative.

Finally, we want to discuss the chiral perturbation theory (χ PT) Wilson coefficients of Refs. [9, 10], based on the present results¹. It is of great interest to study them within the present model, since the model is chiral symmetric by construction. In Refs. [9, 10], it was shown that the non-local quark bilinear operators for the twist-3 meson DAs and their moments are reconstructed in terms of the chiral field, $\Sigma = \exp(i\sqrt{2}\mathcal{M}^a\lambda^a/F_{\mathcal{M}})$, to the leading order (LO) and next-to-leading order (NLO) as follows:

$$\langle \xi^m \rangle_{\mathcal{M}} = \langle 0 | c_m \mathcal{O}_{\text{LO}} + b_m \mathcal{O}_{\text{NLO}} | \mathcal{M}(P) \rangle, \quad (18)$$

n	$a_{2,\pi}^p$	$a_{4,\pi}^p$	$a_{6,\pi}^p$	$a_{1,K}^p$	$a_{2,K}^p$	$a_{3,K}^p$	$a_{4,K}^p$	$a_{5,K}^p$	$a_{6,K}^p$
1	0.44296	0.15081	0.08421	0.02033	0.13715	0.00170	0.10748	-0.00842	0.05975
2	-0.18451	-0.45274	-0.26330	0.02492	-0.43993	-0.02802	-0.32704	-0.01548	-0.16701
3	-0.43067	-0.55590	-0.17839	0.02364	-0.64675	-0.03674	-0.37241	-0.01999	-0.09403
[6]	0.51578	0.25454	0.21624	...	0.26310	...	-0.05216	...	0.14697
[11]	0.43726	-0.07150	-0.19686	0.18372	0.27067	0.39530	-0.24693	0.05497	-0.24361
n	$a_{2,\pi}^\sigma$	$a_{4,\pi}^\sigma$	$a_{6,\pi}^\sigma$	$a_{1,K}^\sigma$	$a_{2,K}^\sigma$	$a_{3,K}^\sigma$	$a_{4,K}^\sigma$	$a_{5,K}^\sigma$	$a_{6,K}^\sigma$
1	0.01658	0.00401	0.00230	-0.00359	-0.00735	-0.00367	0.00368	-0.00280	0.00188
2	-0.09841	-0.01923	-0.00368	-0.00474	-0.11797	-0.00298	-0.01314	-0.00068	-0.00282
3	-0.14770	-0.01596	-0.00052	-0.00466	-0.16072	-0.00335	-0.00949	-0.00002	0.00002
[6]	0.09400	-0.00057	-0.00078	...	0.05203	...	-0.00048	...	-0.00066
[11]	0.09787	-0.00156	-0.00114	0.08976	0.05383	0.05111	-0.01501	0.00734	-0.00578

TABLE I: Gegenbauer moments for the twist-3 pion and kaon distribution amplitudes, $a_{m,\mathcal{M}}^{p,\sigma}$.

¹ The analysis of the χ PT Wilson coefficients can be performed by using the derivative expansion within the present scheme. The corresponding work will be found elsewhere.

where c_m and b_m are the χ PT Wilson coefficients encoding physics in the scale of $\Lambda_{\chi\text{PT}}$. From Eq. (18), one can derive the equations concerning the even and odd moments:

$$\begin{aligned}
\langle \xi^{2m+1} \rangle_\pi^{S,T} &= 0, \\
\langle \xi^{2m+1} \rangle_K^{S,T} &= (m_s - \bar{m})b_{2m+1,2}, \\
\langle \xi^{2m} \rangle_\pi^{S,T} &= \langle \xi^{2m} \rangle_{m_q=0}^{S,T} + 2\bar{m}\alpha_{2m} + (2\bar{m} + m_s)\beta_{2m}, \\
\langle \xi^{2m} \rangle_K^{S,T} &= \langle \xi^{2m} \rangle_{m_q=0}^{S,T} + (\bar{m} + m_s)\alpha_{2m} + (2\bar{m} + m_s)\beta_{2m}
\end{aligned} \tag{19}$$

where β_{2m} and α_{2m} are the collective parameters of c_m and b_m . Note that these equations are model-independent, since they are only based on chiral symmetry of the operators [9, 10]. The first equation in Eq. (19) indicates that the odd moments are all zero for the pion DAs, which is true because of isospin symmetry. The second equation represents flavor SU(3) symmetry breaking effects in terms of the mass difference between the light and strange current quarks. $\bar{m}$ is defined as $(m_u + m_d)/2 = 5$ MeV. As for the even moments, the third and fourth equations are given. They can be simplified due to the fact that $\phi_{\mathcal{M}}^p$ and $\phi_{\mathcal{M}}^\sigma$ are all zero in the chiral limit as easily verified in Eqs. (10) and (11).

$$\alpha_{2m} = \frac{\langle \xi^{2m} \rangle_\pi - \langle \xi^{2m} \rangle_K}{\bar{m} + m_s}, \tag{20}$$

$$\beta_{2m} = \frac{(\bar{m} + m_s)\langle \xi^{2m} \rangle_\pi - 2\bar{m}\langle \xi^{2m} \rangle_K}{m_s^2 + \bar{m}m_s - 2\bar{m}^2}. \tag{21}$$

It is straightforward to obtain the values of b_{2m+1} by using the results listed in Table II. The corresponding results are listed in Table III. Since the b_{2m+1} are the expansion coefficients for the operators of NLO [9, 10], we find that the effects from the NLO are getting weaker as m increases. In addition to b_{2m+1} , we also list the calculated values of α_{2m} in Table III. We verify that the values of β_{2m} are rather tiny (10^{-3}). Note that β_{2m} are the coefficients for the NLO. It turns out that the values of the α_{2m} for the $\phi_{\mathcal{M}}^p$ are rather sizeable, while they are less than 10^{-2} for the $\phi_{\mathcal{M}}^\sigma$.

n	$\langle \xi^2 \rangle_\pi^p$	$\langle \xi^4 \rangle_\pi^p$	$\langle \xi^6 \rangle_\pi^p$	$\langle \xi \rangle_K^p$	$\langle \xi^2 \rangle_K^p$	$\langle \xi^3 \rangle_K^p$	$\langle \xi^4 \rangle_K^p$	$\langle \xi^5 \rangle_K^p$
1	0.39240	0.25445	0.19072	0.00678	0.35162	0.00416	0.21840	0.00292
2	0.30873	0.16742	0.10820	0.00831	0.27468	0.00338	0.14142	0.00160
3	0.27591	0.13666	0.08164	0.00788	0.24710	0.00263	0.11663	0.00081
[7]	0.340 ~ 0.359	0.164 ~ 0.211	...	...	...	...	...	...
[8]	0.52 ± 0.03	0.44 ± 0.01	...	-0.10 ± 0.03	0.43 ± 0.04	...	...	...
[11]	0.38653	0.24510	0.17879	0.06124	0.36757	0.05933	0.22355	0.05198
n	$\langle \xi^2 \rangle_\pi^\sigma$	$\langle \xi^4 \rangle_\pi^\sigma$	$\langle \xi^6 \rangle_\pi^\sigma$	$\langle \xi \rangle_K^\sigma$	$\langle \xi^2 \rangle_K^\sigma$	$\langle \xi^3 \rangle_K^\sigma$	$\langle \xi^4 \rangle_K^\sigma$	$\langle \xi^5 \rangle_K^\sigma$
1	0.20568	0.08992	-0.00186	-0.00216	0.19748	-0.00162	0.08442	-0.00131
2	0.16626	0.06122	-0.00145	-0.00284	0.15955	-0.00179	0.05738	-0.00123
3	0.14936	0.05030	-0.00117	-0.00280	0.14490	-0.00184	0.04799	-0.00125
[8]	0.34 ± 0.03	0.20 ± 0.01	...	-0.13 ± 0.04	0.173 ± 0.002	...	...	...
[11]	0.23250	0.10747	0.06240	0.06124	0.20218	0.03282	0.08949	0.02208

TABLE II: Moments for the twist-3 pion and kaon distribution amplitudes, $\langle \xi^m \rangle_{\mathcal{M}}^{p,\sigma}$.

n	b_1^p	b_3^p	b_5^p	b_1^σ	b_3^σ	b_5^σ
1	0.04675	0.02872	0.02011	-0.01487	-0.01119	-0.00900
2	0.05730	0.02333	0.01105	-0.01961	-0.01231	-0.00849
3	0.05433	0.01812	0.00561	-0.01929	-0.01266	-0.00860
n	α_2^p	α_4^p	α_6^p	α_2^σ	α_4^σ	α_6^σ
1	0.26310	0.23258	0.19845	0.05290	0.03548	-0.31503
2	0.21968	0.16774	0.12555	0.04329	0.02477	-0.18723
3	0.18587	0.12923	0.08890	0.02877	0.01490	-0.14587

TABLE III: χ PT Wilson coefficients b_{2m+1} and α_{2m} defined in Eqs. (19) and (20).

IV. SUMMARY AND CONCLUSION

We investigated the two types of two-particle twist-3 pion and kaon distribution amplitudes, based on the nonlocal chiral quark model from the instanton vacuum. We considered explicitly flavor SU(3) symmetry breaking effects, employing the modified effective chiral action [16, 17]. The current-quark mass dependence was also taken into account for the dynamical quark mass $M_f(k)$ by solving the integral equation for the quark propagator under the instanton effects. We employed simple-pole type parameterization for the form factor in $M_f(k)$.

Numerical results of $\phi_{\mathcal{M}}^p$ and $\phi_{\mathcal{M}}^\sigma$ were given for various cases. We obtained the symmetric pion distribution amplitudes due to isospin symmetry. On the contrary, the kaon distribution amplitudes turned out to be asymmetric because of SU(3) symmetry breaking effects. However, calculated curves for the kaon DAs were almost symmetric in spite of the quark mass differences.

When power $n = 1$ for the dynamical quark mass was used, we found that the results of the $\phi_{\mathcal{M}}^p$ seem to be very similar to those of Ref. [11] with the same end-point behavior. However, as n increased, the results look similar to those of Ref. [7]. As for the $\phi_{\mathcal{M}}^\sigma$, the results were very similar to that of the asymptotic one, with $n = 1$ used. As the power n increases, the kaon distribution amplitudes were suppressed at the end points and were getting narrower. We also investigated the Gegenbauer moments and the moments of the distribution amplitudes.

Finally, we estimated the χ PT Wilson coefficients for the twist-3 pseudoscalar meson distribution amplitudes, using the present results. These estimations may be useful in analyzing the higher-twist meson distribution amplitudes in terms of the chiral field operators. A more systematic analysis can be carried out for the χ PT Wilson coefficients, the derivative expansion being employed. The corresponding analysis will appear soon.

The investigation on the three-particle twist-3 and twist-4 distribution amplitudes with gluon operators is under way.

Acknowledgments

The present work has been supported by the two-year project of Pusan National University. The work of S.i.N. is supported by the Brain Korea 21 (BK21) project in Center of Excellency for Developing Physics Researchers of Pusan National University, Korea. The

authors would like to thank M. M. Musakhanov for his comments on this work. S.i.N. is grateful to Y. Kwon for fruitful discussions.

Appendix

A. Twist-3 pseudoscalar type pseudoscalar meson DA

The twist-3 pseudoscalar type DA can be written by using Eq. (1) as follows:

$$\begin{aligned}\phi_{\mathcal{M}}^p(u) &= \frac{(P \cdot \hat{n})(m_f + m_g)}{m_{\mathcal{M}}^2 f_{\mathcal{M}}} \int_{-\infty}^{\infty} \frac{d\tau}{\pi} e^{-i(2u-1)\tau P \cdot \hat{n}} \langle 0 | \bar{\psi}_f(\tau \hat{n}) i \gamma_5 \psi_g(-\tau \hat{n}) | \mathcal{M}(P) \rangle \\ &= \frac{\sqrt{2}(P \cdot \hat{n})(m_f + m_g)}{m_{\mathcal{M}}^2 F_{\mathcal{M}}} \int_{-\infty}^{\infty} \frac{d\tau}{\pi} e^{-i(2u-1)\tau P \cdot \hat{n}} \langle 0 | \bar{\psi}_f(\tau \hat{n}) i \gamma_5 \psi_g(-\tau \hat{n}) | \mathcal{M}(P) \rangle,\end{aligned}\quad (22)$$

where $F_{\mathcal{M}} = f_{\mathcal{M}}/\sqrt{2}$ stands for the empirical value of the pseudoscalar meson decay constant. We set $m_{\mathcal{M}}$ to be the mass of the pseudoscalar meson. After integrating over τ and k_+ , we arrive at Eq. (10). The trace shown in Eq. (10) can be evaluated in the present framework as follows:

$$\begin{aligned}\text{tr}_d \left[\frac{\sqrt{M_f(k)}}{D(k)} \gamma_5 \frac{\sqrt{M_g(k-P)}}{D(k-P)} \gamma_5 \right] &= 4M_0 \sqrt{f(m_f)f(m_g)} \left[\frac{n\Lambda^2}{n\Lambda^2 - k^2} \right]^n \left[\frac{n\Lambda^2}{n\Lambda^2 - (k-P)^2} \right]^n \\ &\times \left\{ k^2 - k \cdot P - \left[m_f + M_0 f(m_f) \left[\frac{n\Lambda^2}{n\Lambda^2 - k^2} \right]^{2n} \right] \left[m_g + M_0 f(m_g) \left[\frac{n\Lambda^2}{n\Lambda^2 - (k-P)^2} \right]^{2n} \right] \right\} \\ &\times \left\{ k^2 - m_f^2 - 2M_0 f(m_f) \left[\frac{n\Lambda^2}{n\Lambda^2 - k^2} \right]^{2n} - M_0^2 f^2 \left[\frac{n\Lambda^2}{n\Lambda^2 - k^2} \right]^{4n} \right\}^{-1} \\ &\times \left\{ (k-P)^2 - m_g^2 - 2M_0 f(m_g) \left[\frac{n\Lambda^2}{n\Lambda^2 - (k-P)^2} \right]^{2n} - M_0^2 f^2 \left[\frac{n\Lambda^2}{n\Lambda^2 - (k-P)^2} \right]^{4n} \right\}^{-1} \\ &= 4\sqrt{\eta_f \eta_g} \frac{(\beta k_- - k_T^2)(\alpha k_- - \gamma_f)^{3n}(\beta k_- - \gamma_g)^{3n}}{\mathcal{D}_f \mathcal{D}_g} \\ &- 4\sqrt{\eta_f \eta_g} (\alpha k_- + \gamma_f)^n (\beta k_- - \gamma_g)^n \frac{[m_f(\alpha k_- - \gamma_f)^{2n} + \eta_f][m_g(\beta k_- - \gamma_g)^{2n} + \eta_g]}{\mathcal{D}_f \mathcal{D}_g},\end{aligned}\quad (23)$$

where we use the following abbreviations:

$$\begin{aligned}\alpha &= P_+, \quad \beta = (u-1)P_+, \\ \gamma_f &= k_T^2 + n\Lambda^2, \quad \gamma_g = (u-1)m_{\mathcal{M}}^2 + k_T^2 + n\Lambda^2, \\ \delta_f &= k_T^2 + m_f^2, \quad \delta_g = (u-1)m_{\mathcal{M}}^2 + k_T^2 + m_g^2, \\ \eta_f &= M_0 f(m_f)(n\Lambda^2)^{2n}, \quad \eta_g = M_0 f(m_g)(n\Lambda^2)^{2n}.\end{aligned}\quad (24)$$

$\mathcal{D}_{f,g}$ in the denominator of Eq. (23) reads:

$$\mathcal{D}_{f,g} = [(\alpha k_- - \delta_{f,g})(\alpha k_- - \gamma_{f,g})^{4n} - 2m_{f,g}\eta_{f,g}(\alpha k_- - \gamma_{f,g})^{2n} - \eta_{f,g}^2 + i\epsilon]. \quad (25)$$

Finally, we have the following expression for the $\phi_{\mathcal{M}}^p(u)$:

$$\phi_{\mathcal{M}}^p(u) = -\frac{4iN_c\sqrt{\eta_f\eta_g}(m_f + m_g)P_+}{m_{\mathcal{M}}^2 F_{\mathcal{M}}^2} \int \frac{dk_- dk_T^2}{(2\pi)^3} \left[\frac{(\beta k_- - k_T^2)(\alpha k_- - \gamma_f)^{3n}(\beta k_- - \gamma_g)^{3n}}{\mathcal{D}_f \mathcal{D}_g} \right]$$

$$-(\alpha k_- + \gamma_f)^n (\beta k_- - \gamma_g)^n \frac{[m_f(\alpha k_- - \gamma_f)^{2n} + \eta_f][m_g(\beta k_- - \gamma_g)^{2n} + \eta_g]}{\mathcal{D}_f \mathcal{D}_g} \Big]. \quad (26)$$

B. Twist-3 pseudotensor type pseudoscalar meson DA

One can write the pseudotensor type twist-3 DA, using Eq. (2) as follows:

$$\begin{aligned} \phi_{\mathcal{M}}^\sigma(u) = & -\frac{6\sqrt{2}(m_f + m_g)}{m_{\mathcal{M}}^2 F_{\mathcal{M}}} \int_{-\infty}^{\infty} \frac{d\tau}{\pi} \\ & \times \int_0^u dv e^{-2iv\tau P \cdot \hat{n}} \langle 0 | \bar{\psi}_f(\tau \hat{n}) i(P \not{\hat{n}} - P \cdot \hat{n}) \gamma_5 \psi_g(-\tau \hat{n}) | \mathcal{M}(P) \rangle. \end{aligned} \quad (27)$$

The trace in the matrix element of Eq. (27) can be derived as in the case of the pseudoscalar-type one. Having performed the trace, we finally arrive at

$$\phi_{\mathcal{M}}^\sigma(u) = \frac{12iN_c(m_f + m_g)P_+^2}{m_{\mathcal{M}}^2 F_{\mathcal{M}} F_\pi} \int_0^u dv \int \frac{dk_- d^2 k_T}{(2\pi)^2} \frac{k_- (\alpha k_- - \gamma_f)^{3n} (\beta k_- - \gamma_g)^{3n}}{\mathcal{D}_f \mathcal{D}_g}. \quad (28)$$

-
- [1] A. V. Efremov and A. V. Radyushkin, Phys. Lett. B **94**, 245 (1980).
 - [2] G. P. Lepage and S. J. Brodsky, Phys. Lett. B **87**, 359 (1979).
 - [3] G. P. Lepage and S. J. Brodsky, Phys. Rev. D **22**, 2157 (1980).
 - [4] V. L. Chernyak and A. R. Zhitnitsky, Phys. Rept. **112**, 173 (1984).
 - [5] V. M. Braun and I. E. Filyanov, Z. Phys. C **44**, 157 (1989) [Sov. J. Nucl. Phys. **50**, 511.1989 YAFIA,50,818 (1989 YAFIA,50,818-830.1989)].
 - [6] P. Ball, JHEP **9901**, 010 (1999).
 - [7] T. Huang, X. H. Wu and M. Z. Zhou, Phys. Rev. D **70**, 014013 (2004).
 - [8] T. Huang, M. Z. Zhou and X. H. Wu, Eur. Phys. J. C **42**, 271 (2005).
 - [9] J. W. Chen and I. W. Stewart, Phys. Rev. Lett. **92**, 202001 (2004).
 - [10] J. W. Chen, H. M. Tsai and K. C. Weng, Phys. Rev. D **73**, 054010 (2006).
 - [11] P. Ball, V. M. Braun and A. Lenz, JHEP **0605**, 004 (2006).
 - [12] E. V. Shuryak, Nucl. Phys. B **203**, 93 (1982).
 - [13] D. Diakonov and V. Y. Petrov, Nucl. Phys. B **245**, 259 (1984).
 - [14] D. Diakonov and V. Y. Petrov, Nucl. Phys. B **272**, 457 (1986).
 - [15] M. Musakhanov, Eur. Phys. J. C **9**, 235 (1999).
 - [16] M. Musakhanov, hep-ph/0104163.
 - [17] M. Musakhanov, Nucl. Phys. A **699**, 340 (2002).
 - [18] V. Y. Petrov, M. V. Polyakov, R. Ruskov, C. Weiss and K. Goeke, Phys. Rev. D **59**, 114018 (1999).
 - [19] A. E. Dorokhov, JETP Lett. **77**, 63 (2003) [Pisma Zh. Eksp. Teor. Fiz. **77**, 68 (2003)].
 - [20] M. Praszalowicz and A. Rostworowski, Phys. Rev. D **64**, 074003 (2001).
 - [21] M. Praszalowicz and A. Rostworowski, Phys. Rev. D **66**, 054002 (2002).
 - [22] S. i. Nam, H. -Ch. Kim, A. Hosaka and M. M. Musakhanov, Phys. Rev. D **74**, 014019 (2006).
 - [23] P. V. Pobylitsa, Phys. Lett. B **226**, 387 (1989).

- [24] D. Diakonov, M. V. Polyakov and C. Weiss, Nucl. Phys. B **461** (1996) 539.
- [25] S. i. Nam and H. -Ch. Kim, arXiv:hep-ph/0605041.